

Inclass Exercise 12
Geostrophic Flow in the Atmosphere and the Ocean
100 points (Due Friday 4 May 2018)

Part I. You are provided the 500 mb analysis (without data plots) for 12 UTC 9 February 2005. Note the station in southeastern Oklahoma (indicated by the square). The line segment shown is a portion of the "n" axis (normal to the flow) at that location, with the segment length shown there (Δn) = 300 km.

The geostrophic wind speed can be calculated from the expression

$$V_g = -g/f \Delta z / \Delta n$$

where g is gravity, 9.8 m s^{-2} , f is the Coriolis parameter, numerically equal to approximately $10 \times 10^{-5} \text{ s}^{-1}$ at around the latitude of the station shown, Δz is the difference in 500 mb heights normal to the flow measured along the increment Δn (remember, to estimate a gradient, take the value at location 2, furthest along the positive direction of the given axis, and subtract from it the value at location 1.)

1. Estimate the value of the geostrophic wind in meters per second, kilometers per hour and knots at the location shown. Lay out all steps on a separate sheet, show all expressions. $1 \text{ knot} = 0.514 \text{ m s}^{-1}$

Caution, Caution: *the challenge here will be to keep units consistent initially. Please remember that when you solve for the geostrophic wind speed using the above expression, your result will initially be in meters per second. You will need to convert that to kilometers per hour and knots. Remember that a nautical mile is 6040 feet.*

2. Plot the value with conventional weather map symbols for wind direction and wind speed (knots) on the map.
3. With a long arrow/streamline drawn right on the chart, locate the position of the fastest current (the polar jet stream).




Fig. 1: 500 mb chart for 1200 UTC 9 February 2005

Part II. You are provided with the time-averaged topography of the ocean (cm) relative to mean sealevel for the period 1992–2002. Note the two locations at 40N in the eastern and western Pacific. The line segment shown is a portion of the "n" axis (normal to the flow) at that location, with the segment length shown there $\Delta n = 300$ km.

The speed of the geostrophic current can be calculated from the expression

$$V_g = -g/f \Delta z / \Delta n$$

where g is gravity, 9.8 m s^{-2} , f is the Coriolis parameter, numerically equal to approximately $10 \times 10^{-5} \text{ s}^{-1}$ at 40N, Δz is the difference in sea-surface height topography normal to the flow measured along the increment Δn (remember, to estimate a gradient, take the value at location 2, furthest along the positive direction of the given axis, and subtract from it the value at location 1.) In this case, $\Delta n = 300$ km.

1. Estimate the value of the geostrophic current meters per second, kilometers per hour and knots at the locations shown. Lay out all steps, show all expressions.
 $1 \text{ knot} = 0.514 \text{ m s}^{-1}$

Caution, Caution: *the challenge here will be to keep units consistent initially. Please remember that when you solve for the geostrophic wind speed using the above expression, your result will initially be in meters per second. You will need to convert that to kilometers per hour and knots. Remember that a nautical mile is 6040 feet.*

2. With a long arrow/streamline drawn right on Fig. 2, locate the position of the fastest current north of the Tropic of Cancer.




Fig 2: Time-averaged topography of the ocean (cm) relative to mean sealevel for the period 1992–2002

